



Arachis Pintoï

Arachis pintoï

LEGUME · PERENNIAL · TROPICAL / FROST-FREE

PRODUCT OVERVIEW

A persistent, shade-tolerant creeping legume for humid pastures, orchards and perennial-crop groundcover.

WHY SHORTLIST THIS PRODUCT

- Living groundcover**
 Can protect orchard floors when competition, mowing and moisture are actively managed.
- Pasture performance**
 Supports productive grazing systems when entry, residual, rest and fertility are managed as one program.
- Shade adaptation**
 Maintains useful groundcover under partial shade where establishment conditions are suitable.
- Biological nitrogen**
 As a legume, it can support biological nitrogen fixation when compatible rhizobia and sound establishment are used.

PRODUCT SNAPSHOT

CROP TYPE

Legume · Perennial

SEASON / CLIMATE

Tropical / frost-free

PRIMARY ROLES

Orchards · Pasture · Shade

GENERAL SOIL FIT

Confirm locally

SOIL pH

4.5–7.0 · species reference

Establish well. Manage the whole cycle

ESTABLISHMENT

- 01 Choose the window**
Match planting to the product's season, frost risk and the field's moisture outlook.
- 02 Test and prepare**
Use a current soil test, control competing vegetation and prepare the seedbed or planting zone for good contact.
- 03 Calibrate the lot**
Confirm the current seed lot, germination, treatment and whether the recommended rate is bulk seed or pure live seed.
- 04 Establish carefully**
Follow the technical reference for placement and method, then verify emergence and stand uniformity early.
- 05 Support nodulation**
Use the compatible inoculant where required and protect it from heat, drying and incompatible seed treatments.

TECHNICAL REFERENCE

Soil pH
4.5–7.0 · species reference

Planting depth
2–6 cm

Dry matter
8–12 t/ha/year

Crude protein
17–20%

Digestibility
67–71%

Rainfall
>1,100 mm preferred

MANAGEMENT

- 01 Set the objective**
Manage the stand for its primary role—cover, soil recovery, forage, grain, tubers or a planned combination.
- 02 Watch the stage**
Use crop stage and field condition—not the calendar alone—to time grazing, harvest or termination.
- 03 Maintain function**
Monitor moisture, fertility, weeds, pests and stand density; adjust before lost ground becomes difficult to recover.
- 04 Plan what follows**
Coordinate residue, regrowth or termination with the needs and planting window of the next enterprise.
- 05 Verify locally**
Adapt the program to local climate, soils, regulations, animal class and product labeling.

Drought tolerance
Moderate once established

Nitrogen
Up to 300 kg/ha/year

Seed-in-pod rate
10–15 kg/ha

Full cover
≈6 months

Growth habit
Creeping, stoloniferous

Flood tolerance
Moderate

IMPORTANT MANAGEMENT NOTE

Reference values are not performance guarantees. Confirm the current seed lot, label, local agronomic recommendation and any animal-use restrictions before planting or feeding.

Technical basis: Tropical Seeds — Arachis pintoi technical brochure

FIELD NOTE

Rates and performance vary with environment, seed lot and management. Scan for the current online profile and supporting resources.

SCAN FOR UPDATES

