

Ubon Stylo

Stylosanthes guianensis

LEGUME · PERENNIAL · WARM SEASON

PRODUCT OVERVIEW

A four-line tropical forage-legume blend for free-draining, low-fertility acid soils and compatible grass pastures.

WHY SHORTLIST THIS PRODUCT

- Pasture performance**
 Supports productive grazing systems when entry, residual, rest and fertility are managed as one program.
- Managed forage**
 Can support forage goals when harvest stage, animal class and local feeding guidance are considered together.
- Acid-soil adaptation**
 Provides an option for acidic soils when fertility constraints are diagnosed and corrected appropriately.
- Biological nitrogen**
 As a legume, it can support biological nitrogen fixation when compatible rhizobia and sound establishment are used.

PRODUCT SNAPSHOT

CROP TYPE

Legume · Perennial

SEASON / CLIMATE

Warm season

PRIMARY ROLES

Pasture · Forage · Acid soils

GENERAL SOIL FIT

Confirm locally

SOIL pH

4.0–5.0 adapted

Establish well. Manage the whole cycle

ESTABLISHMENT

- 01 Choose the window**
Match planting to the product's season, frost risk and the field's moisture outlook.
- 02 Test and prepare**
Use a current soil test, control competing vegetation and prepare the seedbed or planting zone for good contact.
- 03 Calibrate the lot**
Confirm the current seed lot, germination, treatment and whether the recommended rate is bulk seed or pure live seed.
- 04 Establish carefully**
Follow the technical reference for placement and method, then verify emergence and stand uniformity early.
- 05 Support nodulation**
Use the compatible inoculant where required and protect it from heat, drying and incompatible seed treatments.

TECHNICAL REFERENCE

Soil pH
4.0–5.0 adapted

Pure stand rate
6–8 kg/ha

Planting depth
Surface to 1–2 cm

Cut interval
50–60 days

Dry matter
13–18 t/ha/year

Crude protein
14–20%

MANAGEMENT

- 01 Set the objective**
Manage the stand for its primary role—cover, soil recovery, forage, grain, tubers or a planned combination.
- 02 Watch the stage**
Use crop stage and field condition—not the calendar alone—to time grazing, harvest or termination.
- 03 Maintain function**
Monitor moisture, fertility, weeds, pests and stand density; adjust before lost ground becomes difficult to recover.
- 04 Plan what follows**
Coordinate residue, regrowth or termination with the needs and planting window of the next enterprise.
- 05 Verify locally**
Adapt the program to local climate, soils, regulations, animal class and product labeling.

Rainfall
800–1,600 mm

Residual
15–20 cm

Grass mix rate
2 kg + 8 kg grass/ha

Growth habit
Perennial shrub legume

Cut height
50–60 cm target

High-rainfall limit
Avoid >2,000 mm

IMPORTANT MANAGEMENT NOTE

Reference values are not performance guarantees. Confirm the current seed lot, label, local agronomic recommendation and any animal-use restrictions before planting or feeding.

Technical basis: Tropical Seeds — Ubon Stylo / Stylosanthes guianensis brochure

FIELD NOTE

Rates and performance vary with environment, seed lot and management. Scan for the current online profile and supporting resources.

SCAN FOR UPDATES

